

On Regular Locally Scalar Representations of Graph $\tilde{D}_4$ in Hilbert Spaces

S. A. Kruglyak [†], L. A. Nazarova, A. V. Roiter.


[†] Institute of Mathematics of National Academy of Sciences of Ukraine,
Tereshchenkivska str., 3, Kiev, Ukraine, ind. 01601
E-mail: red@imath.kiev.ua

Representations of quivers corresponding to extended Dynkin graphs are described up to equivalence in [1]. Locally scalar representations of graphs in the category of Hilbert spaces were introduced in [2], and such representations are naturally classified up to unitary equivalence.

Representations of $*$ -algebras generated by linearly related orthogonal projections are studied in [3–10] and others. The connection between locally scalar representations of several graphs (trees, which include also Dynkin graphs) and representations of such $*$ -algebras is stated in [11], and we further use this connection.

The present paper is dedicated to the classification of indecomposable regular (see [12]) locally scalar representations of the graph $\tilde{D}_4$ (for $\tilde{D}_4$ those are indecomposable locally scalar representations in the dimension $(2; 1, 1, 1, 1)$). The answer obtained for the corresponding $*$ -algebra in [5, 6], in our opinion, cannot be satisfactory and definitive. We will obtain explicit formulas, expressing matrix elements of a representation by a character (see [2]) of a locally scalar representation and two “free” real parameters.

1. Let $\mathcal{H}$ be the category of finite-dimensional Hilbert spaces, and $\tilde{D}_4$ be an extended Dynkin graph



Remind (see [2]) that a representation Π of the graph $\tilde{D}_4$ associates a space $H_i \in \mathcal{H}$ to each vertex i ($i = \overline{0, 4}$), and a pair of interadjoint linear operators $\Pi(\gamma_i) = \{\Gamma_{i0}; \Gamma_{0i}\}$ to each edge γ_i , where $\Gamma_{0i} : H_i \rightarrow H_0$, $\Gamma_{i0} = \Gamma_{0i}^*$.

A morphism $C : \Pi \rightarrow \tilde{\Pi}$ is a family $C = \{C_i\}_{i=\overline{0,4}}$ of operators $C_i : \Pi(i) \rightarrow \tilde{\Pi}(i)$ such that

the diagrams

$$\begin{array}{ccc} H_i & \xrightarrow{\Gamma_{ji}} & H_j \\ C_i \downarrow & & \downarrow C_j \\ \tilde{H}_i & \xrightarrow{\tilde{\Gamma}_{ji}} & \tilde{H}_j \end{array}$$

are commutative, i. e. $C_j \Gamma_{ji} = \tilde{\Gamma}_{ji} C_i$.

Let M_i be the set of vertices connected with a vertex i by an edge, $A_i = \sum_{j \in M_i} \Gamma_{ij} \Gamma_{ji}$. Representation Π is called locally scalar [2] if all operators A_i are scalar; $A_i = \alpha_i I_{H_i}$, where I_{H_i} is identity operator in a space H_i . Since A_i is a positive operator, $\alpha_i \geq 0$. A vector $\{\dim \Pi(i)\}$ is a dimension of a finite-dimensional representation Π_i ; if $A_i = f(i) I_{H_i}$ then $\{f(i)\}$ is called a character of locally scalar representation Π .

Further we will denote as $\text{Rep}(\tilde{D}_4, f)$ the category of finite-dimensional locally scalar representations of the graph $\tilde{D}_4$ in $\mathcal{H}$ with given character f .

We will assume that $\alpha_i = f(i) > 0$, $i = \overline{0, 4}$ and the character is normalized: $f(0) = \alpha_0 = 1$.

On the other hand, consider the following $*$ -algebra over the field $\mathbb{C}$:

$$\mathcal{P}_{4,f} = \mathbb{C}\langle p_1, p_2, p_3, p_4 \mid p_i = p_i^* = p_i^2, \sum_{i=1}^4 \alpha_i p_i = e \rangle,$$

where $\alpha_i = f(i)$, e is the identity of the algebra, and the category $\text{Rep } \mathcal{P}_{4,f}$ of finite-dimensional $*$ -representation of the algebra $\mathcal{P}_{4,f}$.

Let $\Pi \in \text{Rep}(\tilde{D}_4, f)$, $\Pi(i) = H_i$, $\pi(\gamma_i) = \{\Gamma_{i,0}; \Gamma_{0,i}\}$. Let us construct a representation π of the algebra $\mathcal{P}_{4,f}$ by the following way: $\pi(p_i) = \frac{1}{\alpha_i} \Gamma_{0,i} \cdot \Gamma_{i,0} = P_i$. If $C : \Pi \rightarrow \tilde{\Pi}$ is a morphism in $\text{Rep}(\tilde{D}_4, f)$ then $C_0 : \pi \rightarrow \tilde{\pi}$ is a morphism in the category $\text{Rep } \mathcal{P}_{4,f}$ (C_0 is the operator interlacing representations π and $\tilde{\pi}$). Define a functor

$$\Phi : \text{Rep}(\tilde{D}_4, f) \rightarrow \text{Rep } \mathcal{P}_{4,f}$$

putting $\Phi(\Pi) = \pi$, $\Phi(C) = C_0$. Clearly, the functor Φ is the equivalence of categories.

Let $\pi \in \text{Rep } \mathcal{P}_{4,f}$ be a representation in the space H_0 . Set $H_i = \text{Im } P_i$, $i = \overline{1, 4}$; $\Gamma_{0,i} : H_i \rightarrow H_0$ is the natural injection of the space H_i into H_0 , then, putting $\Pi(\gamma_i) = \{\Gamma_{0,i}; \Gamma_{0,i}^*\}$, we obtain a representation from $\text{Rep}(\tilde{D}_4, f)$. If $C_0 : \pi \rightarrow \tilde{\pi}$ set $C_i = C_0|_{H_i}$. If $\Phi^{(-1)}(\pi) = \Pi$, $\Phi^{(-1)}(C_0) = \{C_i\}_{i=\overline{0,4}}$ then $\Phi\Phi^{(-1)} \sim I_{\text{Rep } \mathcal{P}_{4,f}}$, $\Phi^{(-1)}\Phi \sim I_{\text{Rep}(\tilde{D}_4, f)}$.

2. Consider representations of the $*$ -algebra $\mathcal{P}_{4,f}$ for

$$0 < \alpha_1 \leq \alpha_2 \leq \alpha_3 \leq \alpha_4 < 1, \sum_{i=1}^4 \alpha_i = 2.$$

(in the other cases representations of the $*$ -algebras $\mathcal{P}_{4,f}$ are reduced to the simplest representations by the Coxeter functors [13]).

Let us make the substitution of generators in the algebra $\mathcal{P}_{4,f}$ [5] :

$$\begin{aligned} x &= \alpha_2 p_2 + \alpha_3 p_3 - \frac{1}{2} \beta_1 e, & \beta_1 &= (2 - \alpha_1 + \alpha_2 + \alpha_3 - \alpha_4)/2, \\ y &= \alpha_1 p_1 + \alpha_3 p_3 - \frac{1}{2} \beta_2 e, & \text{where } \beta_2 &= (2 + \alpha_1 - \alpha_2 + \alpha_3 - \alpha_4)/2, \\ z &= \alpha_1 p_1 + \alpha_2 p_2 - \frac{1}{2} \beta_3 e, & \beta_3 &= (2 + \alpha_1 + \alpha_2 - \alpha_3 - \alpha_4)/2. \end{aligned} \quad (1)$$

Denote also

$$\begin{aligned} \gamma_1 &= (\alpha_1^2 - \alpha_2^2 - \alpha_3^2 + \alpha_4^2)/4, \\ \gamma_2 &= (-\alpha_1^2 + \alpha_2^2 - \alpha_3^2 + \alpha_4^2)/4, \\ \gamma_3 &= (-\alpha_1^2 - \alpha_2^2 + \alpha_3^2 + \alpha_4^2)/4. \end{aligned}$$

It is easy to check that $\gamma_1 \leq \gamma_2 \leq \gamma_3$ and $0 \leq \gamma_2$.

The new generators x, y, z satisfy the system of relations

$$\begin{aligned} \{y, z\} &= \gamma_1 e, \\ \{z, x\} &= \gamma_2 e, & (x + y + z)^2 &= \alpha_4^2 e. \\ \{x, y\} &= \gamma_3 e, \end{aligned} \quad (2)$$

The equalities (1) imply

$$\begin{aligned} p_1 &= \frac{-x + y + z}{2\alpha_1} + \frac{1}{2}e, \\ p_2 &= \frac{x - y + z}{2\alpha_2} + \frac{1}{2}e, \\ p_3 &= \frac{x + y - z}{2\alpha_3} + \frac{1}{2}e, \\ p_4 &= \frac{-x - y - z}{2\alpha_4} + \frac{1}{2}e. \end{aligned} \quad (3)$$

3. Let $\gamma_3 = 0$, then $0 \leq \gamma_2 \leq \gamma_3$ implies $\gamma_2 = 0$, hence $\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = \frac{1}{2}$ (this case is considered in [4]), and so $\gamma_1 = 0$.

In this case the system of relation (2) has a form

$$\begin{aligned} \{y, z\} &= 0, \\ \{z, x\} &= 0, & x^2 + y^2 + z^2 &= \frac{1}{4}e. \\ \{x, y\} &= 0, \end{aligned} \quad (4)$$

Let π be an indecomposable two-dimensional representation of the algebra $\mathcal{P}_{4,f}$ and $\pi(x) = X$, $\pi(y) = Y$, $\pi(z) = Z$.

- a) Let $Z = 0$. The matrix X can be diagonalized as a matrix of self-adjoint operator. The relations of anticommutation imply that the triple $X, Y, Z = 0$ is indecomposable only in the case when the diagonalized matrix X equals $\begin{bmatrix} -\lambda & 0 \\ 0 & \lambda \end{bmatrix}$. Then $Y = \begin{bmatrix} 0 & y_{12} \\ \bar{y}_{12} & 0 \end{bmatrix}$, $y_{12} \neq 0$,

and the element y_{12} can be made positive by the admissible transformations. Therefore, we obtain the case

$$X = \lambda \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}, \quad Y = \mu \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad Z = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}; \quad \lambda > 0, \mu > 0.$$

b) Let $Z \neq 0$ and $X = 0$. Then the matrix Y can be diagonalized: $Y = \begin{bmatrix} -\mu & 0 \\ 0 & \mu \end{bmatrix}$, $\mu > 0$ (in the other cases the triple of matrices turn out to be decomposable). Then from $\{y, z\} = 0$ we obtain $Z = \begin{bmatrix} 0 & z_{12} \\ \bar{z}_{12} & 0 \end{bmatrix}$ and one can reduce the matrix Z (does not changing the Y)

by the admissible transformations to the form $Z = \begin{bmatrix} 0 & i\nu \\ i\nu & 0 \end{bmatrix}$, $\nu > 0$. The triple of matrices X, Y, Z is reduced by means of a unitary matrix $U = \begin{bmatrix} \sqrt{2}/2 & \sqrt{2}/2 \\ -\sqrt{2}/2 & \sqrt{2}/2 \end{bmatrix}$ by the transformation UXU^* , UYU^* , UZU^* to the form

$$X = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \quad Y = \mu \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad Z = \nu \begin{bmatrix} 0 & i \\ -i & 0 \end{bmatrix}; \quad \mu > 0, \nu > 0.$$

c) Let $X \neq 0$, $Z \neq 0$, $Y = 0$. In this case the matrices X, Y, Z can be reduced to the form

$$X = \lambda \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}, \quad Y = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \quad Z = \nu \begin{bmatrix} 0 & i \\ -i & 0 \end{bmatrix}; \quad \lambda > 0, \nu > 0.$$

d) $X \neq 0$, $Y \neq 0$, $Z \neq 0$. In this case the matrices X, Y, Z can be reduced to the form

$$X = \lambda \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}, \quad Y = \mu \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad Z = \nu \begin{bmatrix} 0 & i \\ -i & 0 \end{bmatrix}; \quad \lambda > 0, \mu > 0, \nu \in \mathbb{R}, \nu \neq 0.$$

Thus, unifying these cases, we may consider that

$$X = \lambda \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}, \quad Y = \mu \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad Z = \nu \begin{bmatrix} 0 & i \\ -i & 0 \end{bmatrix},$$

where $\lambda^2 + \mu^2 + \nu^2 = \frac{1}{4}$ (follows from (4)), and

either $\lambda > 0, \mu > 0, \nu \in \mathbb{R}$;

either $\lambda = 0, \mu > 0, \nu > 0$;

or $\lambda > 0, \mu = 0, \nu > 0$.

Formulas (3) imply

$$P_1 = \begin{bmatrix} \frac{1}{2} - \lambda & \mu + \nu i \\ \mu - \nu i & \frac{1}{2} + \lambda \end{bmatrix}, \quad P_2 = \begin{bmatrix} \frac{1}{2} + \lambda & -\mu + \nu i \\ -\mu - \nu i & \frac{1}{2} - \lambda \end{bmatrix},$$

$$P_3 = \begin{bmatrix} \frac{1}{2} + \lambda & \mu - \nu i \\ \mu + \nu i & \frac{1}{2} - \lambda \end{bmatrix}, \quad P_4 = \begin{bmatrix} \frac{1}{2} - \lambda & -\mu - \nu i \\ -\mu + \nu i & \frac{1}{2} + \lambda \end{bmatrix}.$$

If $\mu + \nu i = \sqrt{\mu^2 + \nu^2} e^{i\varphi} = \sqrt{\frac{1}{4} - \lambda^2} e^{i\varphi}$ then, passing on to the unitary equivalent representation by means of the matrix $U = \begin{bmatrix} e^{-i\varphi} & 0 \\ 0 & 1 \end{bmatrix}$, we may consider that

$$\begin{aligned} P_1 &= \begin{bmatrix} \frac{1}{2} - \lambda & \sqrt{\frac{1}{4} - \lambda^2} \\ \sqrt{\frac{1}{4} - \lambda^2} & \frac{1}{2} + \lambda \end{bmatrix}, & P_2 &= \begin{bmatrix} \frac{1}{2} + \lambda & e^{i\chi} \sqrt{\frac{1}{4} - \lambda^2} \\ e^{-i\chi} \sqrt{\frac{1}{4} - \lambda^2} & \frac{1}{2} - \lambda \end{bmatrix}, \\ P_3 &= \begin{bmatrix} \frac{1}{2} + \lambda & -e^{-i\chi} \sqrt{\frac{1}{4} - \lambda^2} \\ -e^{-i\chi} \sqrt{\frac{1}{4} - \lambda^2} & \frac{1}{2} - \lambda \end{bmatrix}, & P_4 &= \begin{bmatrix} \frac{1}{2} - \lambda & -\sqrt{\frac{1}{4} - \lambda^2} \\ -\sqrt{\frac{1}{4} - \lambda^2} & \frac{1}{2} + \lambda \end{bmatrix}, \end{aligned} \quad (5)$$

$$\begin{aligned} 0 &\leq \lambda < 1/2, \text{ and} \\ 0 &< \chi < \pi/2 \text{ when } \lambda = 0, \\ -\pi/2 &< \chi \leq \pi/2 \text{ when } 0 < \lambda < 1/2. \end{aligned}$$

4. Let $\gamma_3 \neq 0$, then $\{x, y\} = \gamma_3 e$ implies $X \neq 0$ and $Y \neq 0$. Moreover, X and Y has no zero eigenvalues. Indeed, let matrix X has the form after diagonalization: $X = \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix}$. In the matrices Y and Z either $y_{12} \neq 0$ or $z_{12} \neq 0$ (or else the triple of matrices is decomposable). $\{x, y\} = \gamma_3 e$, $\{x, z\} = \gamma_2 e$ imply $(\lambda_1 + \lambda_2)z_{12} = 0$ and $(\lambda_1 + \lambda_2)y_{12} = 0$. Therefore we can conclude that $-\lambda_1 = \lambda_2 = \lambda \neq 0$, $\lambda > 0$. The same reasoning is useful also for Y .

Let $X = \begin{bmatrix} -\lambda & 0 \\ 0 & \lambda \end{bmatrix}$. $\{x, y\} = \gamma_3 e$ implies $y_{11} = -\frac{\gamma_3}{2\lambda}$, $y_{22} = \frac{\gamma_3}{2\lambda}$; $\{x, z\} = \gamma_2 e$ implies $z_{11} = -\frac{\gamma_2}{2\lambda}$, $z_{22} = \frac{\gamma_2}{2\lambda}$.

$$X = \lambda \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}, \quad Y = \frac{1}{2\lambda} \begin{bmatrix} -\gamma_3 & y_{12} \\ \bar{y}_{12} & \gamma_3 \end{bmatrix}, \quad Z = \frac{1}{2\lambda} \begin{bmatrix} -\gamma_2 & z_{12} \\ \bar{z}_{12} & \gamma_2 \end{bmatrix}.$$

$\{y, z\} = \gamma_1 e$ implies

$$\frac{1}{4\lambda^2} \begin{bmatrix} 2\gamma_2\gamma_3 + y_{12}\bar{z}_{12} + \bar{y}_{12}z_{12} & 0 \\ 0 & 2\gamma_2\gamma_3 + y_{12}\bar{z}_{12} + \bar{y}_{12}z_{12} \end{bmatrix} = \gamma_1 I,$$

hence

$$y_{12}\bar{z}_{12} + \bar{y}_{12}z_{12} = 4\gamma_1\lambda^2 - 2\gamma_2\gamma_3. \quad (6)$$

Let us turn to the unitary equivalent representation by means of the unitary matrix $U = \begin{bmatrix} e^{i\varphi} & 0 \\ 0 & 0 \end{bmatrix}$ so that $y_{12} + z_{12} = r_1$ would be real positive; at that $-y_{12} + z_{12}$ remains to be complex in general, $-y_{12} + z_{12} = r_2 e^{i\chi}$.

Then (2) implies

$$(X + Y + Z)^2 = \frac{1}{4\lambda^2} \begin{bmatrix} (2\lambda^2 + \gamma_2 + \gamma_3)^2 + (y_{12} + z_{12})^2 & 0 \\ 0 & (2\lambda^2 + \gamma_2 + \gamma_3)^2 + (y_{12} + z_{12})^2 \end{bmatrix} = \alpha_4^2 I$$

and

$$r_1^2 = (y_{12} + z_{12})^2 = 4\alpha_4^2\lambda^2 - (2\lambda^2 + \gamma_2 + \gamma_3)^2,$$

from which it easy to obtain

$$\begin{aligned} r_1 &= \sqrt{-4\lambda^4 + 2(\alpha_1^2 + \alpha_4^2)\lambda^2 - \frac{1}{4}(\alpha_4^2 - \alpha_1^2)^2} = \\ &= \sqrt{-4\left(\lambda^2 - \frac{(\alpha_4^2 - \alpha_1^2)^2}{4}\right)\left(\lambda^2 - \frac{(\alpha_4^2 + \alpha_1^2)^2}{4}\right)}, \\ \frac{\alpha_4 - \alpha_1}{2} &\leq \lambda \leq \frac{\alpha_4 + \alpha_1}{2}. \end{aligned} \tag{7}$$

$$\begin{cases} y_{12} + z_{12} = r_1, \\ -y_{12} + z_{12} = r_2 e^{i\chi}, \end{cases} \quad \text{therefore} \quad \begin{cases} y_{12} = (r_1 - r_2 e^{i\chi})/2, \\ z_{12} = (r_1 + r_2 e^{i\chi})/2, \end{cases}$$

$$y_{12}\bar{z}_{12} + \bar{y}_{12}z_{12} = (r_1^2 - r_2^2)/2 = 4\gamma_1\lambda^2 - 2\gamma_2\gamma_3 \text{ (the last equality follows from (6)).}$$

Thus,

$$r_2^2 = r_1^2 + 4\gamma_2\gamma_3 - 8\gamma_1\lambda^2 = -4\lambda^4 + 2(\alpha_2^2 + \alpha_3^2)\lambda^2 - \frac{1}{4}(\alpha_3^2 - \alpha_2^2)^2 \geq 0$$

and

$$\begin{aligned} -y_{12} + z_{12} = r_2 e^{i\chi} &= e^{i\chi} \sqrt{-4\lambda^4 + 2(\alpha_2^2 + \alpha_3^2)\lambda^2 - \frac{1}{4}(\alpha_3^2 - \alpha_2^2)^2} = \\ &= e^{i\chi} \sqrt{-4\left(\lambda^2 - \frac{(\alpha_3^2 - \alpha_2^2)^2}{4}\right)\left(\lambda^2 - \frac{(\alpha_3^2 + \alpha_2^2)^2}{4}\right)}, \\ \frac{\alpha_3 - \alpha_2}{2} &\leq \lambda \leq \frac{\alpha_3 + \alpha_2}{2}. \end{aligned}$$

Now we can directly pass on to the determining of the operators of the representation P_1, P_2, P_3, P_4 :

$$\begin{aligned} P_1 &= \frac{-X+Y+Z}{2\alpha_1} + \frac{1}{2}I = \\ &= \frac{1}{4\alpha_1\lambda} \left[\frac{2\lambda^2 + 2\alpha_1\lambda - \frac{1}{2}(\alpha_4^2 - \alpha_1^2)}{\sqrt{-4\lambda^4 + 2(\alpha_1^2 + \alpha_4^2)\lambda^2 - \frac{1}{4}(\alpha_4^2 - \alpha_1^2)^2}} \frac{\sqrt{-4\lambda^4 + 2(\alpha_1^2 + \alpha_4^2)\lambda^2 - \frac{1}{4}(\alpha_4^2 - \alpha_1^2)^2}}{-2\lambda^2 + 2\alpha_1\lambda + \frac{1}{2}(\alpha_4^2 - \alpha_1^2)} \right] = \\ &= \frac{1}{4\alpha_1\lambda} \left[\frac{2\left(\lambda - \frac{1}{2}(\alpha_4 - \alpha_1)\right)\left(\lambda + \frac{1}{2}(\alpha_4 + \alpha_1)\right)}{\sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_4 - \alpha_1)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_4 + \alpha_1)^2\right)}} \frac{\sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_4 - \alpha_1)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_4 + \alpha_1)^2\right)}}{-2\left(\lambda + \frac{1}{2}(\alpha_4 - \alpha_1)\right)\left(\lambda - \frac{1}{2}(\alpha_4 + \alpha_1)\right)} \right], \end{aligned}$$

$$\begin{aligned} P_2 &= \frac{X-Y+Z}{2\alpha_2} + \frac{1}{2}I = \\ &= \frac{1}{4\alpha_2\lambda} \left[\frac{-2\lambda^2 + 2\alpha_2\lambda + \frac{1}{2}(\alpha_3^2 - \alpha_2^2)}{e^{-i\chi} \sqrt{-4\lambda^4 + 2(\alpha_2^2 + \alpha_3^2)\lambda^2 - \frac{1}{4}(\alpha_3^2 - \alpha_2^2)^2}} \frac{e^{i\chi} \sqrt{-4\lambda^4 + 2(\alpha_2^2 + \alpha_3^2)\lambda^2 - \frac{1}{4}(\alpha_3^2 - \alpha_2^2)^2}}{2\lambda^2 + 2\alpha_2\lambda - \frac{1}{2}(\alpha_3^2 - \alpha_2^2)} \right] = \\ &= \frac{1}{4\alpha_2\lambda} \left[\frac{-2\left(\lambda - \frac{1}{2}(\alpha_3 + \alpha_2)\right)\left(\lambda + \frac{1}{2}(\alpha_3 - \alpha_2)\right)}{e^{-i\chi} \sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_3 + \alpha_2)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_3 - \alpha_2)^2\right)}} \frac{e^{i\chi} \sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_3 + \alpha_2)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_3 - \alpha_2)^2\right)}}{2\left(\lambda + \frac{1}{2}(\alpha_3 + \alpha_2)\right)\left(\lambda - \frac{1}{2}(\alpha_3 - \alpha_2)\right)} \right], \end{aligned}$$

$$\begin{aligned}
P_3 &= \frac{X+Y-Z}{2\alpha_3} + \frac{1}{2}I = \\
&= \frac{1}{4\alpha_3\lambda} \left[\frac{-2\lambda^2 + 2\alpha_3\lambda - \frac{1}{2}(\alpha_3^2 - \alpha_2^2)}{-e^{-i\chi} \sqrt{-4\lambda^4 + 2(\alpha_2^2 + \alpha_3^2)\lambda^2 - \frac{1}{4}(\alpha_3^2 - \alpha_2^2)^2}} \quad -e^{-i\chi} \sqrt{-4\lambda^4 + 2(\alpha_2^2 + \alpha_3^2)\lambda^2 - \frac{1}{4}(\alpha_3^2 - \alpha_2^2)^2}} \right] = \\
&= \frac{1}{4\alpha_3\lambda} \left[\frac{-2\left(\lambda - \frac{1}{2}(\alpha_3 + \alpha_2)\right)\left(\lambda - \frac{1}{2}(\alpha_3 - \alpha_2)\right)}{-e^{-i\chi} \sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_3 + \alpha_2)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_3 - \alpha_2)^2\right)}} \quad -e^{-i\chi} \sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_3 + \alpha_2)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_3 - \alpha_2)^2\right)}} \right], \\
P_4 &= \frac{-X-Y-Z}{2\alpha_4} + \frac{1}{2}I = \\
&= \frac{1}{4\alpha_4\lambda} \left[\frac{2\lambda^2 + 2\alpha_4\lambda + \frac{1}{2}(\alpha_4^2 - \alpha_1^2)}{-\sqrt{-4\lambda^4 + 2(\alpha_4^2 + \alpha_1^2)\lambda^2 - \frac{1}{4}(\alpha_4^2 - \alpha_1^2)^2}} \quad -\sqrt{-4\lambda^4 + 2(\alpha_4^2 + \alpha_1^2)\lambda^2 - \frac{1}{4}(\alpha_4^2 - \alpha_1^2)^2}} \right] = \\
&= \frac{1}{4\alpha_4\lambda} \left[\frac{2\left(\lambda + \frac{1}{2}(\alpha_4 + \alpha_1)\right)\left(\lambda + \frac{1}{2}(\alpha_4 - \alpha_1)\right)}{-\sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_4 + \alpha_1)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_4 - \alpha_1)^2\right)}} \quad -\sqrt{-4\left(\lambda^2 - \frac{1}{4}(\alpha_4 + \alpha_1)^2\right)\left(\lambda^2 - \frac{1}{4}(\alpha_4 - \alpha_1)^2\right)}} \right].
\end{aligned}$$

5. Let us construct, according to p. 1, isometries $\Gamma_{0,i} : H_i \rightarrow H_0$, $i = \overline{1,4}$ by projectors $P_i : H_0 \rightarrow H_0$ and obtain regular indecomposable locally scalar representations of the graph $\tilde{D}_4$:

$$\begin{aligned}
\Gamma_{0,1} &= \begin{bmatrix} \sqrt{\frac{\lambda^2 + \alpha_1\lambda - (\alpha_4^2 - \alpha_1^2)/4}{2\alpha_1\lambda}} \\ \sqrt{\frac{-\lambda^2 + \alpha_1\lambda + (\alpha_4^2 - \alpha_1^2)/4}{2\alpha_1\lambda}} \end{bmatrix}, \quad \Gamma_{0,2} = \begin{bmatrix} \sqrt{\frac{-\lambda^2 + \alpha_2\lambda + (\alpha_3^2 - \alpha_2^2)/4}{2\alpha_2\lambda}} \\ e^{-i\chi} \sqrt{\frac{\lambda^2 + \alpha_2\lambda - (\alpha_3^2 - \alpha_2^2)/4}{2\alpha_2\lambda}} \end{bmatrix}, \\
\Gamma_{0,3} &= \begin{bmatrix} \sqrt{\frac{-\lambda^2 + \alpha_3\lambda - (\alpha_3^2 - \alpha_2^2)/4}{2\alpha_3\lambda}} \\ -e^{-i\chi} \sqrt{\frac{\lambda^2 + \alpha_3\lambda + (\alpha_3^2 - \alpha_2^2)/4}{2\alpha_3\lambda}} \end{bmatrix}, \quad \Gamma_{0,4} = \begin{bmatrix} \sqrt{\frac{\lambda^2 + \alpha_4\lambda + (\alpha_4^2 - \alpha_1^2)/4}{2\alpha_4\lambda}} \\ -\sqrt{\frac{-\lambda^2 + \alpha_4\lambda - (\alpha_4^2 - \alpha_1^2)/4}{2\alpha_4\lambda}} \end{bmatrix}. \\
&0 < (\alpha_4 - \alpha_1)/2 \leq \lambda \leq \min((\alpha_2 + \alpha_3)/2, (\alpha_1 + \alpha_4)/2), \quad -\pi < \chi \leq \pi.
\end{aligned}$$

Thus, indecomposable regular (not necessarily normalized) locally scalar representations of the graph $\tilde{D}_4$ depend on 6 real parameters (on the 4 of 5 parameters $\alpha'_1, \alpha'_2, \alpha'_3, \alpha'_4, \alpha'_0$, connected by the relation $\alpha'_1 + \alpha'_2 + \alpha'_3 + \alpha'_4 = 2\alpha'_0$, $\alpha_i = \frac{\alpha'_i}{\alpha'_0}$, and parameters λ and χ).

References

1. *L. A. Nazarova* Representations of quivers of infinite type. (Russian) *Izv. Akad. Nauk SSSR Ser. Mat.* 37 (1973), 752–791.

2. *S. A. Kruglyak, A. V. Roiter* Locally scalar representations of graphs in the category of Hilbert spaces. *Funct. Anal. Appl.* 39 (2005), no. 2, 91–105
3. *Yu. N. Bespalov*. Sets of orthoprojectors connected by relations. *Ukrainian Math. J.* 44 (1992), no. 3, 269–277.
4. *V. Ostrovskiy and Yu. Samoilenko* Introduction to the theory of representations of finitely presented $*$ -algebras. I. Representations by bounded operators. *Reviews in Mathematics and Mathematical Physics*, 11, pt.1. Harwood Academic Publishers, Amsterdam, 1999. iv+261 pp. ISBN: 90-5823-042-2
5. *D. V. Galinsky, S. A. Kruglyak* Representations of $*$ -algebras generated by linearly related orthogonal projections. (Ukrainian.) *Visnyk Kyiv. derzh. Univ.* No. 2, 24–31, 1999.
6. *D. V. Galinsky*. Representations of $*$ -algebras generated by orthogonal projections satisfying a linear relation. *Methods Funct. Anal. Topology* 4 (1998), no. 3, 27–32.
7. *D. V. Galinsky, M. A. Muratov* On representations of algebras generated by sets of three and four orthoprojections. *Spectral and evolutionary problems*, Vol. 8 (Sevastopol, 1997), 15–22, Tavria Publ., Simferopol, 1998.
8. *S. A. Kruglyak, V. I. Rabanovich, Yu. S. Samoilenko* On sums of projections. *Funct. Anal. Appl.* 36 (2002), no. 3, 182–195
9. *S. Kruglyak, A. Kyrychenko*. On four orthogonal projections that satisfy the linear relation $\alpha_1 P_1 + \alpha_2 P_2 + \alpha_3 P_3 + \alpha_4 P_4 = I$, $\alpha_i > 0$. *Symmetry in nonlinear mathematical physics*, Part 1, 2 (Kyiv, 2001), 461–465, *Pr. Inst. Mat. Nats. Akad. Nauk Ukr. Mat. Zastos.*, 43, Part 1, 2, *Natsional. Akad. Nauk Ukraïni, Inst. Mat.*, Kiev, 2002.
10. *A. Strelets*. Description of the representations of the algebras generated by four linearly related idempotents. *J. Algebra Appl.* 4 (2005), no. 6, 671–681.
11. *S. Kruglyak, S. Popovich, Yu. Samoilenko*. The spectral problem and $*$ -representations of algebras associated with Dynkin graphs. *J. Algebra Appl.* 4 (2005), no. 6, 761–776.
12. *I. K. Redchuk, A. V. Roiter*. Singular locally scalar representations of quivers in Hilbert spaces, and separating functions. *Ukrainian Math. J.* 56 (2004), no. 6, 947–963.
13. *S. A. Kruglyak*. Coxeter functors for a certain class of $*$ -quivers and $*$ -algebras. *Methods Funct. Anal. Topology* 8 (2002), no. 4, 49–57.